





In summary, an acceptable daily intake (ADI) of 200 micrograms per day and an occupational exposure limit of 20 micrograms/m<sup>3</sup> are recommended for methylpredisolone succinate.

**References:**

1. US National Cancer Institute, NCI Drug Dictionary, <http://www.cancer.gov/Templates/drugdictionary.aspx?CdrID=41528>
2. RxList, ■■■■■■■■■■■■■■■■■■■■■■ for injection, USP, <http://www.rxlist.com/cgi/generic/solumed.htm>.
3. Japanese Pharmacopoeia (JP15) Database, ■■■■■■■■■■■■■■■■■■■■■■, [http://molddb.nihs.go.jp/jp/scripts/name\\_exact\\_eng.asp](http://molddb.nihs.go.jp/jp/scripts/name_exact_eng.asp).
4. ScienceLab.com, Material Safety Data Sheet for 10/10/2005.
5. Pfizer Inc., Prescribing Information for ■■■■■■■■■■■■■■■■■■■■■■ for injection, USP, June 2006.
6. American Society of Health System Pharmacists, ■■■■■■■■■■■■■■■■■■■■■■, [http://www.ashp.org/s\\_ashp/bulletin.asp](http://www.ashp.org/s_ashp/bulletin.asp). April 8, 2008.
7. Heron, RJL and FC Pickering, (2003), Health effects of exposure to active pharmaceutical ingredients (APIs), Occupational Medicine, 53:357-362.
8. Newton RW, MCK Browning, J Iqbal, N Piercy, and DG Adamson, (1978), Adrenocortical suppression in workers manufacturing synthetic glucocorticoids, British Medical Journal, 1:73-74.
9. Pezzarossa, A., et.al., (1987), Adrenal suppression in workers employed in manufacturing synthetic glucocorticoids: solutions to a problem, Lancet, 1:515.
10. Newton RW, MCK Browning, J Iqbal, N Piercy, and DG Adamson, (1982), Adrenocortical suppression in workers manufacturing synthetic glucocorticoids: solutions to a problem, British Medical Journal, 39:179-182.
11. US National Library of Medicine, Hazardous Substances Database, ■■■■■■■■■■■■■■■■■■■■■■ ■■■■■■■■■■■■■■■■■■■■■■, <http://toxnet.nlm.nih.gov>.

Respectfully submitted,

Edward V. Sargent MPH PhD DABT  
Managing Director  
EV Sargent LLC  
1430 Gulf Boulevard  
Suite 205  
Clearwater, Florida 33767  
USA  
Date Prepared: June 10, 2008